

MAP1LC3A Rabbit mAb

Catalog No.: A12319 **Recombinant** **5 Publications**

Basic Information

Observed MW

14kDa,16kDa/16kDa

Calculated MW

14kDa

Category

Primary antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC2636

Background

MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. The protein encoded by this gene is one of the light chain subunits and can associate with either MAP1A or MAP1B. Two transcript variants encoding different isoforms have been found for this gene. The expression of variant 1 is suppressed in many tumor cell lines, suggesting that may be involved in carcinogenesis.

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID

84557

Swiss Prot

Q9H492

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human MAP1LC3A (Q9H492).

Synonyms

LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3; MAP1LC3A

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Immunoprecipitation analysis of 300 µg extracts of NIH/3T3 cells using 3 µg MAP1LC3A antibody (A12319). Western blot was performed from the immunoprecipitate using MAP1LC3A antibody (A12319) at a dilution of 1:1000. NIH/3T3 cells were treated by Chloroquine (50 µM) at 37°C for 20 hours.



Western blot analysis of lysates from SH-SY5Y cells using MAP1LC3A Rabbit mAb(A12319) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Western blot analysis of lysates from NIH/3T3, C6 cells using MAP1LC3A Rabbit mAb (A12319) at 1:1000 dilution. NIH/3T3 cells and C6 cells were treated by Chloroquine (50 µM) at 37°C for 20 hours.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.