# **CGB Rabbit pAb**

Catalog No.: A12419 1 Publications



## **Basic Information**

#### **Observed MW**

22kDa

#### **Calculated MW**

18kDa

### Category

Primary antibody

## **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene is a member of the glycoprotein hormone beta chain family and encodes the beta 3 subunit of chorionic gonadotropin (CG). Glycoprotein hormones are heterodimers consisting of a common alpha subunit and an unique beta subunit which confers biological specificity. CG is produced by the trophoblastic cells of the placenta and stimulates the ovaries to synthesize the steroids that are essential for the maintenance of pregnancy. The beta subunit of CG is encoded by 6 genes which are arranged in tandem and inverted pairs on chromosome 19q13.3 and contiguous with the luteinizing hormone beta subunit gene.

## **Recommended Dilutions**

**WB** 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID**Swiss Prot
1082
PODN86

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 21-165 of human CGB (NP\_000728.1).

## **Synonyms**

CGB; LHB; CGB5; CGB7; CGB8; hCGB

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates, using CGB Rabbit pAb (A12419) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.