

[KO Validated] CD133 Rabbit pAb

Catalog No.: A12711 **KO Validated** **3 Publications**

Basic Information

Observed MW

120kDa

Calculated MW

97kDa

Category

Primary antibody

Applications

ELISA, WB, IF/ICC, IP

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a pentaspan transmembrane glycoprotein. The protein localizes to membrane protrusions and is often expressed on adult stem cells, where it is thought to function in maintaining stem cell properties by suppressing differentiation. Mutations in this gene have been shown to result in retinitis pigmentosa and Stargardt disease. Expression of this gene is also associated with several types of cancer. This gene is expressed from at least five alternative promoters that are expressed in a tissue-dependent manner. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:1000**IF/ICC** 1:50 - 1:200**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID

8842

Swiss Prot

O43490

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 179-433 of human CD133 (NP_006008.1).

Synonyms

RP41; AC133; CD133; MCDR2; STGD4; CORD12; PROML1; MSTP061; 33

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data



Immunoprecipitation analysis of 200 µg extracts of HT-29 cells using 3 µg CD133 antibody (A12711). Western blot was performed from the immunoprecipitate using CD133 antibody (A12711) at a dilution of 1:1000.



Western blot analysis of lysates from wild type (WT) and CD133 knockout (KO) HCT116 cells, using [KO Validated] CD133 Rabbit pAb (A12711) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



Western blot analysis of lysates from HCT116 cells, using [KO Validated] CD133 Rabbit pAb (A12711) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.