# MAFF Rabbit pAb

Catalog No.: A12920



### **Basic Information**

#### **Observed MW**

18kDa

#### **Calculated MW**

18kDa

#### Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human

### **Background**

The protein encoded by this gene is a basic leucine zipper (bZIP) transcription factor that lacks a transactivation domain. It is known to bind the US-2 DNA element in the promoter of the oxytocin receptor (OTR) gene and most likely heterodimerizes with other leucine zippercontaining proteins to enhance expression of the OTR gene during term pregnancy. The encoded protein can also form homodimers, and since it lacks a transactivation domain, the homodimer may act as a repressor of transcription. This gene may also be involved in the cellular stress response. Multiple transcript variants encoding two different isoforms have been found for this gene.

### **Recommended Dilutions**

**WB** 

1:500 - 1:2000

### **Immunogen Information**

**Gene ID** 23764

**Swiss Prot** 

Q9ULX9

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 75-164 of human MAFF (NP\_036455.1).

### **Synonyms**

U-MAF; hMafF; MAFF

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from HT-29 cells, using MAFF Rabbit pAb (A12920) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.