

Cyclin D1 Rabbit pAb

Catalog No.: A1301SP

32 Publications

Basic Information

Observed MW

36 kDa

Calculated MW

34 kDa

Category

Primary antibody

Applications

WB, IP, IF/ICC, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with tumor suppressor protein Rb and the expression of this gene is regulated positively by Rb. Mutations, amplification and overexpression of this gene, which alters cell cycle progression, are observed frequently in a variety of human cancers.

Recommended Dilutions

WB 1:5000 - 1:10000**IP** 0.5 µg - 4 µg antibody for
200 µg - 400 µg extracts of
whole cells**IF/ICC** 1:400 - 1:1400**IHC-P** 1:2000 - 1:8000**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on your
specific assay
requirements. For high-ratio
antibody dilutions
(≥1:10000) a sequential
dilution method is strongly
recommended to ensure
measurement accuracy.

Immunogen Information

Gene ID

595

Swiss Prot

P24385

Immunogen

This information is considered to be commercially sensitive.

Synonyms

BCL1; PRAD1; U21B31; D11S287E; Cyclin D1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide.

May contain 0.05% BSA as specified on the Certificate of Analysis.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Immunoprecipitation of Cyclin D1 from 300 µg extracts of HeLa cells was performed using 2 µg of Cyclin D1 Rabbit pAb (A1301SP). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Cyclin D1 Rabbit pAb (A1301SP) at a dilution of 1:5000.




Western blot analysis of various lysates using Cyclin D1 Rabbit pAb (A1301SP) at 1:6000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5 s.




Confocal imaging of MCF7 cells using Cyclin D1 Rabbit pAb (A1301SP, dilution 1:1300) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of RAW 264.7 cells using Cyclin D1 Rabbit pAb (A1301SP, dilution 1:1300) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of PC-12 cells using Cyclin D1 Rabbit pAb (A1301SP, dilution 1:1300) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using Cyclin D1 Rabbit pAb (A1301SP) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using Cyclin D1 Rabbit pAb (A1301SP) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human mantle cell lymphoma tissue using Cyclin D1 Rabbit pAb (A1301SP) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using Cyclin D1 Rabbit pAb (A1301SP) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat skin tissue using Cyclin D1 Rabbit pAb (A1301SP) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.