# LRPAP1 Rabbit pAb

Catalog No.: A13026



### **Basic Information**

#### **Observed MW**

42kDa

### **Calculated MW**

41kDa

### Category

Primary antibody

### **Applications**

ELISA, WB, IHC-P, IF/ICC

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a protein that interacts with the low density lipoprotein (LDL) receptorrelated protein and facilitates its proper folding and localization by preventing the binding of ligands. Mutations in this gene have been identified in individuals with myopia 23. Alternative splicing results in multiple transcript variants.

### **Recommended Dilutions**

WB	1:500 - 1:2000
IHC-P	1:100 - 1:200
IF/ICC	1:50 - 1:200

## **Immunogen Information**

Gene ID	Swiss Prot	
4043	P30533	

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 35-357 of human LRPAP1 (NP\_002328.1).

### **Synonyms**

RAP; MRAP; A2RAP; HBP44; MYP23; A2MRAP; alpha-2-MRAP; LRPAP1

### Contact

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using LRPAP1 Rabbit pAb (A13026) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunofluorescence analysis of U-2 OS cells using LRPAP1 Rabbit pAb (A13026) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.