# TTC5 Rabbit pAb

Catalog No.: A13232



### **Basic Information**

#### **Observed MW**

49kDa

### **Calculated MW**

49kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Predicted to enable DNA binding activity and chromatin binding activity. Predicted to be involved in DNA repair. Predicted to act upstream of or within positive regulation of transcription by RNA polymerase II. Located in nucleoplasm.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 91875

**Swiss Prot** 

Q8N0Z6

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human TTC5 (NP\_612385.2).

# **Synonyms**

Strap; NEDCAFD; TTC5

# **Contact**

| <b>a</b>  |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| $\odot$   | T | www.abclonal.com.cn       |


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using TTC5 Rabbit pAb (A13232) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.