# **GARS Rabbit pAb**

Catalog No.: A13305



## **Basic Information**

#### **Observed MW**

70-83kDa

### **Calculated MW**

83kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IF/ICC

### **Cross-Reactivity**

Human, Mouse

## **Background**

This gene encodes glycyl-tRNA synthetase, one of the aminoacyl-tRNA synthetases that charge tRNAs with their cognate amino acids. The encoded enzyme is an (alpha)2 dimer which belongs to the class II family of tRNA synthetases. It has been shown to be a target of autoantibodies in the human autoimmune diseases, polymyositis or dermatomyositis. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

## **Immunogen Information**

**Gene ID Swiss Prot**2617
P41250

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 60-370 of human GARS (NP\_002038.2).

### **Synonyms**

GARS; HMN5; CMT2D; DSMAV; GlyRS; HMN5A; SMAD1; SMAJI

### **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using GARS antibody (A13305) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.