

# PELP1 Rabbit pAb

Catalog No.: A13414 **1 Publications**

## Basic Information

### Observed MW

**Calculated MW**  
120kDa

**Category**  
Primary antibody

**Applications**  
ELISA, IHC-P

**Cross-Reactivity**  
Human, Mouse, Rat

## Background

This gene encodes a transcription factor which coactivates transcription of estrogen receptor responsive genes and corepresses genes activated by other hormone receptors or sequence-specific transcription factors. Expression of this gene is regulated by both members of the estrogen receptor family. This gene may be involved in the progression of several types of cancer. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

IHC-P 1:50 - 1:200

## Immunogen Information

<b>Gene ID</b> 27043	<b>Swiss Prot</b> Q8IZL8
-------------------------	-----------------------------

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 884-1130 of human PELP1 (Q8IZL8).

### Synonyms

MNAR; P160; PELP1

## Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

## Product Information


<b>Source</b> Rabbit	<b>Isotype</b> IgG	<b>Purification</b> Affinity purification
-------------------------	-----------------------	--

### Storage


Store at -20°C. Avoid freeze / thaw cycles.  
Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Validation Data


---



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using PELP1 Rabbit pAb (A13414) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using PELP1 Rabbit pAb (A13414) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen using PELP1 Rabbit pAb (A13414) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.