Smac/Diablo Rabbit pAb

Catalog No.: A13432



Basic Information

Observed MW

Calculated MW 27kDa

Category

Primary antibody

Applications

IF/ICC,ELISA

Cross-Reactivity

Human

Background

This gene encodes an inhibitor of apoptosis protein (IAP)-binding protein. The encoded mitochondrial protein enters the cytosol when cells undergo apoptosis, and allows activation of caspases by binding to inhibitor of apoptosis proteins. Overexpression of the encoded protein sensitizes tumor cells to apoptosis. A mutation in this gene is associated with young-adult onset of nonsyndromic deafness-64. Alternative splicing results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

IF/ICC 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID56616

Swiss Prot
Q9NR28

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

SMAC; DFNA64; Smac/Diablo

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\overline{a}	ī	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Immunofluorescence analysis of MCF-7 cells using Smac/Diablo Rabbit pAb (A13432). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.