# ICAM-1/CD54 Rabbit pAb

Catalog No.: A13620



### **Basic Information**

#### **Observed MW**

92kDa

#### **Calculated MW**

58kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# **Immunogen Information**

Gene ID

Swiss Prot

3383

P05362

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to a mino acids 250-480 of human ICAM-1/CD54 (NP\_000192.2).

### **Synonyms**

BB2; CD54; P3.58; ICAM-1/CD54

### **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of Raji, using ICAM-1/CD54 antibody (A13620) at 1:800 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.