# **THOC7 Rabbit pAb**

Catalog No.: A13700 1 Publications



## **Basic Information**

#### **Observed MW**

24kDa

#### **Calculated MW**

24kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Predicted to enable RNA binding activity. Involved in mRNA export from nucleus and viral mRNA export from host cell nucleus. Located in cytosol and nuclear speck. Part of THO complex part of transcription export complex. Colocalizes with chromosome, telomeric region.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 80145

**Swiss Prot** 

Q6I9Y2

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-204 of human THOC7 (NP\_079351.2).

### **Synonyms**

fSAP24; hTREX30; NIF3L1BP1; THOC7

### **Contact**

| 2         |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| •         | T | www.abclonal.com.cn       |


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using THOC7 Rabbit pAb (A13700) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.