

ASCC2 Rabbit pAb

Catalog No.: A13789

Basic Information

Observed MW

110kDa

Calculated MW

86kDa

Category

Primary antibody

Applications

ELISA, WB

Cross-Reactivity

Human, Mouse, Rat

Background

Predicted to enable ubiquitin binding activity. Involved in regulation of transcription, DNA-templated; rescue of stalled ribosome; and ribosome-associated ubiquitin-dependent protein catabolic process. Located in nucleus. Part of activating signal cointegrator 1 complex.

Recommended Dilutions

WB 1:500 - 1:2000

Immunogen Information

Gene ID

84164

Swiss Prot

Q9H118

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 488-757 of human ASCC2 (NP_115580.2).

Synonyms

p100; ASC1p100; ASCC2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using ASCC2 Rabbit pAb (A13789) at 1:3000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.