Leader in Biomolecular Solutions for Life Science

# **IKKe Rabbit pAb**

Catalog No.: A14093



### **Basic Information**

Observed MW 80kDa

Calculated MW 80kDa

**Category** Primary antibody

Applications WB,IHC-P,ELISA

Cross-Reactivity Human, Mouse, Rat

### Background

IKBKE is a noncanonical I-kappa-B (see MIM 164008) kinase (IKK) that is essential for regulating antiviral signaling pathways. IKBKE has also been identified as a breast cancer (MIM 114480) oncogene and is amplified and overexpressed in over 30% of breast carcinomas and breast cancer cell lines (Hutti et al., 2009 [PubMed 19481526]).

### **Recommended Dilutions**

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Immunogen Information**

#### **Gene ID** 9641

Swiss Prot Q14164

#### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### Synonyms

IKKE; IKKI; IKK-E; IKK-i; IKKE

### Contact


6	400-999-6126
$\mathbf{X}$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification** Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using IKK $\epsilon$  Rabbit pAb (A14093) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunohistochemistry analysis of paraffinembedded Rat spleen using IKKɛ Rabbit pAb (A14093) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse brain using IKKɛ Rabbit pAb (A14093) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.