# **HSD17B6** Rabbit pAb

Catalog No.: A14262



### **Basic Information**

#### **Observed MW**

36kDa

### **Calculated MW**

36kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Mouse

# **Background**

The protein encoded by this gene has both oxidoreductase and epimerase activities and is involved in androgen catabolism. The oxidoreductase activity can convert 3 alpha-adiol to dihydrotestosterone, while the epimerase activity can convert androsterone to epiandrosterone. Both reactions use NAD+ as the preferred cofactor. This gene is a member of the retinol dehydrogenase family.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# **Immunogen Information**

Gene ID 8630 Swiss Prot 014756

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 178-317 of human  $HSD17B6\ (NP\_003716.2)$ .

### **Synonyms**

HSE; RODH; SDR9C6; HSD17B6

### **Contact**

| 6         |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| •         | T | www.abclonal.com.cn       |

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using HSD17B6 Rabbit pAb (A14262) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.