# F8A1 Rabbit pAb

Catalog No.: A14806



### **Basic Information**

#### **Observed MW**

39kDa

### **Calculated MW**

39kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is conserved in the mouse implies it has some function. Unlike factor VIII, this gene is transcribed abundantly in a wide variety of cell types.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID 8263 **Swiss Prot** 

P23610

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 200-300 of human F8A1 ( $NP_036283.2$ ).

# **Synonyms**

F8A; HAP40; DXS522E; F8A1

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using F8A1 Rabbit pAb (A14806) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.