AIF1/IBA1 Rabbit pAb

Catalog No.: A1527 12 Publications



Basic Information

Observed MW

17kDa

Calculated MW

17kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human

Background

This gene encodes a protein that binds actin and calcium. This gene is induced by cytokines and interferon and may promote macrophage activation and growth of vascular smooth muscle cells and T-lymphocytes. Polymorphisms in this gene may be associated with systemic sclerosis. Alternative splicing results in multiple transcript variants, but the full-length and coding nature of some of these variants is not certain.

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 μg/mL. Please optimize the

concentration based on your specific assay requirements.

Immunogen Information

Gene IDSwiss Prot
199
P55008

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

IBA1; IRT1; AIF-1; IRT-1; AIF1/IBA1

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	Τ	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using AIF1/IBA1 Rabbit pAb (A1527) at 1:5000 dilution incubated overnight at 4° C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Negative control (NC): HeLa Exposure time: 10s.



Immunohistochemistry analysis of paraffinembedded Human esophageal cancer using AIF1/IBA1 Rabbit pAb (A1527) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human tonsil using AIF1/IBA1 Rabbit pAb (A1527) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.