# DCAF11 Rabbit pAb

Catalog No.: A15519 5 Publications



# **Basic Information**

### **Observed MW**

61kDa

### **Calculated MW**

62kDa

## Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a WD repeat-containing protein that interacts with the COP9 signalosome, a macromolecular complex that interacts with cullin-RING E3 ligases and regulates their activity by hydrolyzing cullin-Nedd8 conjugates. Multiple alternatively spliced transcript variants have been found for this gene.

# **Recommended Dilutions**

**WB** 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**80344

Swiss Prot
Q8TEB1

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-170 of human DCAF11 (NP\_079506.3).

## **Synonyms**

GL014; WDR23; PRO2389; DCAF11

## **Contact**

| <b>a</b>  |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| $\odot$   | Т | www.abclonal.com.cn       |


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from Mouse liver, using DCAF11 Rabbit pAb (A15519) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.