# **NIFK Rabbit pAb**

Catalog No.: A15595 2 Publications



## **Basic Information**

### **Observed MW**

39kDa

### **Calculated MW**

34kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a protein that interacts with the forkhead-associated domain of the Ki-67 antigen. The encoded protein may bind RNA and may play a role in mitosis and cell cycle progression. Multiple pseudogenes exist on chromosomes 5, 10, 12, 15, and 19.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting

concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay

requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
84365
Q9BYG3

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

Nopp34; MKI67IP; NIFK

## Contact

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Т	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using NIFK Rabbit pAb (A15595) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.