# Calnexin Rabbit pAb

Catalog No.: A15631 38 Publications



## **Basic Information**

#### **Observed MW**

90kDa

## **Calculated MW**

68kDa

### Category

Primary antibody

## **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a member of the calnexin family of molecular chaperones. The encoded protein is a calcium-binding, endoplasmic reticulum (ER)-associated protein that interacts transiently with newly synthesized N-linked glycoproteins, facilitating protein folding and assembly. It may also play a central role in the quality control of protein folding by retaining incorrectly folded protein subunits within the ER for degradation. Alternatively spliced transcript variants encoding different isoforms have been described.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$ 

Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID Swiss Prot**821
P27824

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

CNX; P90; IP90; Calnexin

# **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
<u> </u>	ī	www.ahclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using Calnexin Rabbit pAb (A15631) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of PC-12 cells using Calnexin Rabbit pAb (A15631) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.