# NAB1 Rabbit pAb

Catalog No.: A15692 1 Publications



## **Basic Information**

#### **Observed MW**

54kDa

### **Calculated MW**

54kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse

# **Background**

Predicted to enable transcription coregulator activity. Predicted to be involved in regulation of transcription, DNA-templated. Predicted to act upstream of or within endochondral ossification; nervous system development; and regulation of epidermis development. Predicted to be located in nucleoplasm. Predicted to be active in nucleus.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 4664 Q13506

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 189-317 of human NAB1 (NP\_005957.2).

## **Synonyms**

NAB1

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{\triangle}$	ī	www.ahclonal.com.cn


### **Product Information**

**Purification Source** Isotype Rabbit IgG Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of various lysates using NAB1 Rabbit pAb (A15692) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.