# PTPRD Rabbit pAb

Catalog No.: A15713 1 Publications



#### **Basic Information**

#### **Observed MW**

300kDa

#### **Calculated MW**

215kDa

## Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular region, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus represents a receptor-type PTP. The extracellular region of this protein is composed of three lg-like and eight fibronectin type III-like domains. Studies of the similar genes in chicken and fly suggest the role of this PTP is in promoting neurite growth, and regulating neurons axon guidance. Multiple alternatively spliced transcript variants of this gene have been reported. A related pseudogene has been identified on chromosome 5.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

Gene ID 5789 **Swiss Prot** 

P23468

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1100-1265 of human PTPRD (NP\_002830.1).

#### **Synonyms**

HPTP; PTPD; HPTPD; HPTPDELTA; RPTPDELTA; R-PTP-delta; PTPRD

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	1	www.abclonal.com.cn

#### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using PTPRD Rabbit pAb (A15713) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3min.