# TRIM62 Rabbit pAb

Catalog No.: A15855



### **Basic Information**

#### **Observed MW**

54kDa

### **Calculated MW**

54kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IHC-P

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Enables identical protein binding activity; transcription coactivator activity; and ubiquitin-protein transferase activity. Involved in several processes, including negative regulation of viral transcription; positive regulation of NF-kappaB transcription factor activity; and positive regulation of antifungal innate immune response. Is active in cytoplasm.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IHC-P** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**Swiss Prot
55223
Q9BVG3

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 120-280 of human TRIM62 (NP\_060677.2).

## **Synonyms**

DEAR1; TRIM62

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{\triangle}$	ī	www.ahclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using TRIM62 Rabbit pAb (A15855) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunohistochemistry analysis of paraffinembedded Human colon using TRIM62 Rabbit pAb (A15855) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.