FER1L5 Rabbit pAb

Catalog No.: A15926



Basic Information

Observed MW

242kDa

Calculated MW

238kDa

Category

Primary antibody

Applications

ELISA,WB,IF/ICC

Cross-Reactivity

Mouse, Rat

Background

Predicted to enable calcium ion binding activity and calcium-dependent phospholipid binding activity. Predicted to be involved in several processes, including myeloid cell activation involved in immune response; negative regulation of phagocytosis; and plasma membrane organization. Predicted to be located in plasma membrane. Predicted to be integral component of membrane. Predicted to be active in T-tubule and cytoplasmic vesicle.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID90342

Swiss Prot
A0AVI2

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1350-1670 of human FER1L5 (NP_001280012.1).

Synonyms

FER1L5

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\Box}$	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using FER1L5 Rabbit pAb (A15926) at 1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit $\lg G$ (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 150s.



Immunofluorescence analysis of paraffinembedded rat testis using FER1L5 Rabbit pAb (A15926) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffinembedded mouse testis using FER1L5 Rabbit pAb (A15926) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.