# **MAGOHB Rabbit pAb**

Catalog No.: A16192 1 Publications



### **Basic Information**

#### **Observed MW**

17kDa

### **Calculated MW**

17kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Rat

# **Background**

Enables RNA binding activity. Involved in mRNA splicing, via spliceosome and nuclear-transcribed mRNA catabolic process, nonsense-mediated decay. Located in nucleus. Part of U2-type catalytic step 1 spliceosome; U2-type precatalytic spliceosome; and exon-exon junction complex.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 55110

**Swiss Prot** 

Q96A72

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-148 of human MAGOHB (NP\_060518.1).

### **Synonyms**

MGN2; mago; magoh; MAGOHB

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using MAGOHB Rabbit pAb (A16192) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.