

# TriMethyl-Histone H3-K27 Mouse mAb

Catalog No.: A16199 **16 Publications**

## Basic Information

### Observed MW

17 kDa

### Calculated MW

15 kDa

### Category

Primary antibody

### Applications

WB, IHC-P, IF/ICC, DB, ChIP, CUT&Tag, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

AMC0015

## Recommended Dilutions

**WB** 1:2000 - 1:10000

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

**DB** 1:500 - 1:2000

**ChIP** 5µg antibody for  
5µg-10µg of Chromatin

**CUT&Tag** 10<sup>5</sup> cells /1 µg

**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

## Immunogen Information

### Gene ID

8350

### Swiss Prot

P68431

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; TriMethyl-Histone H3-K27

## Product Information

### Source

Mouse

### Isotype

IgG1, kappa

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Contact

---


 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)


 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

---




## Validation Data



Western blot analysis of various lysates using TriMethyl-Histone H3-K27 Mouse mAb (A16199) at 1:10000 dilution incubated overnight at 4°C.  
 Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Negative control (NC): Recombinant H3 protein (10 ng per lane) .  
 Exposure time: 20 s.





Chromatin immunoprecipitation analysis of extracts of HeLa; cells, using TriMethyl-Histone H3-K27 Mouse mAb antibody (A16199) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Dot-blot analysis of all sorts of methylation peptides using TriMethyl-Histone H3-K27 antibody (A16199) at 1:1000 dilution.

Confocal imaging of NIH/3T3 cells using TriMethyl-Histone H3-K27 Mouse mAb (A16199, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.

Confocal imaging of C6 cells using TriMethyl-Histone H3-K27 Mouse mAb (A16199, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina(RK20265)




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using

## Validation Data


---

(A16199, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using TriMethyl-Histone H3-K27 Mouse mAb (A16199) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

from  $10^5$  K562 cells with 1 µg TriMethyl-Histone H3-K27 Mouse mAb antibody (A16199), along with a Goat Anti-Mouse IgG (H+L). The CUT&Tag results indicate the enrichment pattern of TriMethyl-Histone H3-K27 in representative gene loci (MYOD1), as shown in figure.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using TriMethyl-Histone H3-K27 Mouse mAb (A16199) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

TriMethyl-Histone H3-K27 Mouse mAb (A16199) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using TriMethyl-Histone H3-K27 Mouse mAb (A16199) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.