# **USP22 Rabbit pAb**

Catalog No.: A16297 1 Publications



## **Basic Information**

#### **Observed MW**

60kDa

### **Calculated MW**

60kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Rat

## **Background**

Enables enzyme binding activity; nuclear receptor coactivator activity; and thiol-dependent deubiquitinase. Contributes to H4 histone acetyltransferase activity. Involved in positive regulation of mitotic cell cycle; positive regulation of transcription, DNA-templated; and protein modification by small protein conjugation or removal. Part of SAGA complex.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 23326

Swiss Prot

Q9UPT9

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human USP22 (NP\_056091.1).

## **Synonyms**

USP3L; USP22

### **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using USP22 Rabbit pAb (A16297) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.