

NR2F1 Rabbit pAb

Catalog No.: A16437 **1 Publications**

Basic Information

Observed MW

Refer to figures

Calculated MW

46kDa

Category

Primary antibody

Applications

IHC-P,ELISA

Cross-Reactivity

Human

Background

The protein encoded by this gene is a nuclear hormone receptor and transcriptional regulator. The encoded protein acts as a homodimer and binds to 5'-AGGTCA-3' repeats. Defects in this gene are a cause of Bosch-Boonstra optic atrophy syndrome (BBOAS).

Recommended Dilutions

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

7025

Swiss Prot

P10589

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 60-180 of human NR2F1 (NP_005645.1).

Synonyms

EAR3; BBOAS; EAR-3; SVP44; BBSOAS; ERBAL3; COUPTF1; TFCOUP1; COUP-TFI; TCFCOUP1; NR2F1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using NR2F1 Rabbit pAb (A16437) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using NR2F1 Rabbit pAb (A16437) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.