

FOXG1 Rabbit pAb

Catalog No.: A16851 **2 Publications**

Basic Information

Observed MW

52kDa

Calculated MW

52kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This locus encodes a member of the fork-head transcription factor family. The encoded protein, which functions as a transcriptional repressor, is highly expressed in neural tissues during brain development. Mutations at this locus have been associated with Rett syndrome and a diverse spectrum of neurodevelopmental disorders defined as part of the FOXG1 syndrome. This gene is dysregulated in many types of cancer and is the target of multiple microRNAs that regulate the proliferation of tumor cells.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

2290

Swiss Prot

P55316

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

BF1; BF2; QIN; FKH2; HBF2; HFK1; HFK2; HFK3; KHL2; FHKL3; FKHL1; FKHL2; FKHL3; FKHL4; HBF-1; HBF-2; HBF-3; FOXG1A; FOXG1B; FOXG1C; HBF-G2; FOXG1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using FOXG1 Rabbit pAb (A16851) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.