# **DPF3 Rabbit pAb**

Catalog No.: A16890



## **Basic Information**

#### **Observed MW**

43kDa

### **Calculated MW**

43kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Mouse, Rat

## **Background**

This gene encodes a member of the D4 protein family. The encoded protein is a transcription regulator that binds acetylated histones and is a component of the BAF chromatin remodeling complex. Alternate splicing results in multiple transcript variants encoding different isoforms.

## **Recommended Dilutions**

1:500 - 1:1000 **WB** 

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID Swiss Prot** 8110 Q92784

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 90-200 of human DPF3 (NP\_036206.3).

### **Synonyms**

CERD4; BAF45C; SMARCG3; DPF3

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Τ	www.abclonal.com.cn


## **Product Information**

**Purification Source** Isotype Rabbit IgG Affinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from Mouse heart, using DPF3 Rabbit pAb (A16890) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.