# ADRA2C Rabbit pAb

Catalog No.: A16956



## **Basic Information**

#### **Observed MW**

50kDa

#### **Calculated MW**

50kDa

## Category

Primary antibody

## **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse

# **Background**

Alpha-2-adrenergic receptors are members of the G protein-coupled receptor superfamily. They include 3 highly homologous subtypes: alpha2A, alpha2B, and alpha2C. These receptors have a critical role in regulating neurotransmitter release from sympathetic nerves and from adrenergic neurons in the central nervous system. The mouse studies revealed that both the alpha2A and alpha2C subtypes were required for normal presynaptic control of transmitter release from sympathetic nerves in the heart and from central noradrenergic neurons. The alpha2A subtype inhibited transmitter release at high stimulation frequencies, whereas the alpha2C subtype modulated neurotransmission at lower levels of nerve activity. This gene encodes the alpha2C subtype, which contains no introns in either its coding or untranslated sequences.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID**Swiss Prot
152
P18825

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

ADRA2L2; ADRARL2; ADRA2RL2; ALPHA2CAR; ADRA2C

# Contact

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using ADRA2C Rabbit pAb (A16956) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.