

HIF1 β /ARNT Rabbit pAb

Catalog No.: A16990

Basic Information

Observed MW

Refer to figures

Calculated MW

87kDa

Category

Primary antibody

Applications

ELISA,IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a protein containing a basic helix-loop-helix domain and two characteristic PAS domains along with a PAC domain. The encoded protein binds to ligand-bound aryl hydrocarbon receptor and aids in the movement of this complex to the nucleus, where it promotes the expression of genes involved in xenobiotic metabolism. This protein is also a co-factor for transcriptional regulation by hypoxia-inducible factor 1. Chromosomal translocation of this locus with the ETV6 (ets variant 6) gene on chromosome 12 have been described in leukemias. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

405

Swiss Prot

P27540

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 465-570 of human HIF1 β /ARNT (NP_001659.1).

Synonyms

HIF1B; TANGO; bHLHe2; HIF1BETA; HIF-1beta; HIF1-beta; HIF-1-beta; HIF1 β /ARNT

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data



Immunofluorescence analysis of H9C2 cells using HIF1 β /ARNT Rabbit pAb (A16990) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using HIF1 β /ARNT Rabbit pAb (A16990) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-S OS cells using HIF1 β /ARNT Rabbit pAb (A16990) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.