# **AXIN2 Rabbit pAb**

Catalog No.: A17022 1 Publications



## **Basic Information**

#### **Observed MW**

100kDa

### **Calculated MW**

94kDa

#### Category

Primary antibody

## **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

The Axin-related protein, Axin2, presumably plays an important role in the regulation of the stability of beta-catenin in the Wnt signaling pathway, like its rodent homologs, mouse conductin/rat axil. In mouse, conductin organizes a multiprotein complex of APC (adenomatous polyposis of the colon), beta-catenin, glycogen synthase kinase 3-beta, and conductin, which leads to the degradation of beta-catenin. Apparently, the deregulation of beta-catenin is an important event in the genesis of a number of malignancies. The AXIN2 gene has been mapped to 17q23-q24, a region that shows frequent loss of heterozygosity in breast cancer, neuroblastoma, and other tumors. Mutations in this gene have been associated with colorectal cancer with defective mismatch repair.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting concentration is 1  $\mu$ g/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**8313
Swiss Prot
Q9Y2T1

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 400-500 of human AXIN2 (NP\_004646.3).

## **Synonyms**

AXIL; ODCRCS; AXIN2

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from SW480 cells, using AXIN2 Rabbit pAb (A17022) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.