BAG4 Rabbit pAb

Catalog No.: A17052



Basic Information

Observed MW 68kDa

Calculated MW 50kDa

Category Primary antibody

Applications ELISA,WB,IHC-P

Cross-Reactivity Human, Mouse, Rat

Background

The protein encoded by this gene is a member of the BAG1-related protein family. BAG1 is an anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. This protein was found to be associated with the death domain of tumor necrosis factor receptor type 1 (TNF-R1) and death receptor-3 (DR3), and thereby negatively regulates downstream cell death signaling. The regulatory role of this protein in cell death was demonstrated in epithelial cells which undergo apoptosis while integrin mediated matrix contacts are lost. Alternatively spliced transcript variants encoding distinct isoforms have been identified.

Recommended Dilutions

Immunogen Information

 WB
 1:500 - 1:2000
 Gene ID
 Swiss Prot

 9530
 095429

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human BAG4 (NP_001191807.1).

Synonyms

SODD; BAG-4; BAG4

Contact


a 400-999-6126 x cn.market@abclonal.com.cn y www.abclonal.com.cn

Product Information

Source Rabbit **lsotype** IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using BAG4 Rabbit pAb (A17052) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.