# **APOBEC3F Rabbit pAb**

Catalog No.: A17269



### **Basic Information**

#### **Observed MW**

45kDa

#### **Calculated MW**

45kDa

### Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human

### **Background**

This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been identified.

### **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 200316

**Swiss Prot** 

L6 Q8IUX4

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 189-373 of human APOBEC3F (NP\_660341.2).

#### **Synonyms**

A3F; KA6; ARP8; BK150C2.4.MRNA; APOBEC3F

### **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn

#### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from PC3 cells, using APOBEC3F Rabbit pAb (A17269) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.