NR4A2 Rabbit pAb

Catalog No.: A17318



Basic Information

Observed MW

67kDa

Calculated MW

67kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human, Mouse

Background

This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. The encoded protein may act as a transcription factor. Mutations in this gene have been associated with disorders related to dopaminergic dysfunction, including Parkinson disease, schizophernia, and manic depression. Misregulation of this gene may be associated with rheumatoid arthritis. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID 4929 **Swiss Prot**

P43354

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 90-170 of human NR4A2 (NP $_$ 006177.1).

Synonyms

NOT; RNR1; HZF-3; IDLDP; NURR1; TINUR; NR4A2

Contact

2		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.




Western blot analysis of various lysates using (A17318) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Western blot analysis of lysates from Mouse brain, using (A17318) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit $\lg G$ (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.