# CD146/MCAM Rabbit pAb

Catalog No.: A17347



### **Basic Information**

#### **Observed MW**

118kDa

### **Calculated MW**

72kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Rat

# **Background**

Involved in glomerular filtration and vascular wound healing. Acts upstream of or within angiogenesis. Located in external side of plasma membrane. Biomarker of chronic obstructive pulmonary disease and uveal melanoma.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID 4162 **Swiss Prot** 

P43121

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 340-560 of human CD146/CD146/MCAM (NP\_006491.2).

### **Synonyms**

CD146; MUC18; HEMCAM; METCAM; MelCAM; CD146/MCAM

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from NCI-H460 cells, using CD146/CD146/MCAM Rabbit pAb (A17347) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.