Leader in Biomolecular Solutions for Life Science



Catalog No.: A17609 1 Publications



# **Basic Information**

Observed MW 146kDa

Calculated MW 146kDa

Category Primary antibody

Applications ELISA,WB

Cross-Reactivity Human, Mouse

# Background

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes an additional conjugation factor, E4, which is involved in multiubiquitin chain assembly. This gene is also the strongest candidate in the neuroblastoma tumor suppressor genes. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

# **Recommended Dilutions**

1:500 - 1:2000

# **Immunogen Information**

WB

### Gene ID 10277

**Swiss Prot** 095155

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 130-250 of human UBE4B (NP\_006039.2).

### Synonyms

E4; UFD2; HDNB1; UBOX3; UFD2A; UBE4B

| Contac | t |
|--------|---|
|--------|---|


# a 400-999-6126 x cn.market@abclonal.com.cn y www.abclonal.com.cn

# **Product Information**

**Source** Rabbit **Isotype** IgG **Purification** Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using UBE4B Rabbit pAb (A17609) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.