# Tubulin beta-1 chain Rabbit pAb

Catalog No.: A17779



## **Basic Information**

#### **Observed MW**

50kDa

#### **Calculated MW**

50kDa

#### Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Mouse

## **Background**

This gene encodes a member of the beta tubulin protein family. Beta tubulins are one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is specifically expressed in platelets and megakaryocytes and may be involved in proplatelet production and platelet release. A mutations in this gene is associated with autosomal dominant macrothrombocytopenia. Two pseudogenes of this gene are found on chromosome Y.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 81027

**Swiss Prot** 

Q9H4B7

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 362-451 of human Tubulin beta-1 chain (NP\_110400.1).

### **Synonyms**

MACTHC1; Tubulin beta-1 chain

## **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using Tubulin beta-1 chain Rabbit pAb (A17779) at 1:1000 dilution

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.