# CTSK Rabbit pAb

Catalog No.: A1782 22 Publications



## **Basic Information**

#### **Observed MW**

26kDa/37kDa

### **Calculated MW**

37kDa

#### Category

Primary antibody

## **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

The protein encoded by this gene is a lysosomal cysteine proteinase involved in bone remodeling and resorption. This protein, which is a member of the peptidase C1 protein family, is predominantly expressed in osteoclasts. However, the encoded protein is also expressed in a significant fraction of human breast cancers, where it could contribute to tumor invasiveness. Mutations in this gene are the cause of pycnodysostosis, an autosomal recessive disease characterized by osteosclerosis and short stature.

## **Recommended Dilutions**

WB 1:100 - 1:500

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**Swiss Prot
1513
P43235

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

CTSO; PKND; PYCD; CTS02; CTS01; CTS02; CTSK

## **Contact**

	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of various lysates using CTSK Rabbit pAb (A1782) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.