# MT-ND2 Rabbit pAb

Catalog No.: A17968 5 Publications



### **Basic Information**

#### **Observed MW**

39kDa

### **Calculated MW**

39kDa

#### Category

Primary antibody

### **Applications**

ELISA, WB, IHC-P, IF/ICC

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Enables NADH dehydrogenase (ubiquinone) activity. Involved in mitochondrial electron transport, NADH to ubiquinone and mitochondrial respiratory chain complex I assembly. Part of mitochondrial respiratory chain complex I. Implicated in Leber hereditary optic neuropathy; multiple sclerosis; myocardial infarction; neurodegenerative disease (multiple); and urinary bladder cancer.

## **Recommended Dilutions**

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

# **Immunogen Information**

Gene ID	Swiss Prot
4536	P03891

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 248-347 of human MT-ND2 ( $YP_003024027.1$ ).

### **Synonyms**

MTND2; ND2; MT-ND2

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.


Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.




Western blot analysis of various lysates using MT-ND2 Rabbit pAb (A17968) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.



Immunohistochemistry analysis of MT-ND2 in paraffin-embedded human colon carcinoma using MT-ND2 Rabbit pAb (A17968) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of HeLa cells using MT-ND2 Rabbit pAb (A17968) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.