# IRF2BP2 Rabbit pAb

Catalog No.: A17985



## **Basic Information**

#### **Observed MW**

61kDa

#### **Calculated MW**

61kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes an interferon regulatory factor-2 (IRF2) binding protein that interacts with the C-terminal transcriptional repression domain of IRF2. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 359948

**Swiss Prot** 

Q7Z5L9

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 475-545 of human IRF2BP2 (Q7Z5L9).

### **Synonyms**

CVID14; IRF2BP2

### **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of lysates from THP-1 cells, using IRF2BP2 Rabbit pAb (A17985) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Western blot analysis of lysates from Mouse spleen, using IRF2BP2 Rabbit pAbd (A17985) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.