

Vgll4 Rabbit pAb

Catalog No.: A18248

11 Publications

Basic Information

Observed MW

31kDa

Calculated MW

31kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Predicted to enable transcription coactivator binding activity. Involved in several processes, including negative regulation of cardiac muscle cell proliferation; negative regulation of hippo signaling; and positive regulation of cellular protein catabolic process. Located in nucleus. Is expressed in lung; submandibular gland primordium; tooth; and vibrissa. Orthologous to human VGLL4 (vestigial like family member 4).

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

232334

Swiss Prot

Q80V24

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

VGL-4; Vgll4

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.


Validation Data




Western blot analysis of lysates from OVCAR3 cells, using Vgll4 Rabbit pAb (A18248) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Vgll4 Rabbit pAb (A18248) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using Vgll4 Rabbit pAb (A18248) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using Vgll4 Rabbit pAb (A18248) at dilution of 1:100 (20x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.