Leader in Biomolecular Solutions for Life Science

PCIF1 Rabbit pAb

Catalog No.: A18347



Basic Information

Observed MW 80kDa

Calculated MW 81kDa

Category Primary antibody

Applications ELISA,WB,IHC-P

Cross-Reactivity Human, Mouse, Rat

Background

Enables RNA polymerase II C-terminal domain phosphoserine binding activity; S-adenosyl-Lmethionine binding activity; and mRNA (2'-O-methyladenosine-N6-)-methyltransferase activity. Involved in mRNA methylation; negative regulation of translation; and positive regulation of translation. Located in intercellular bridge; microtubule cytoskeleton; and nucleoplasm.

Recommended Dilutions

Immunogen Information

WB	1:1000 - 1:5000	Gene ID	Swiss Prot	
IHC-P	1:50 - 1:200	63935	Q9H4Z3	

Immunogen

Recombinant fusion protein containing a sequence corresponding to a mino acids 1-200 of human PCIF1 (NP_071387.1).

Synonyms

CAPAM; hCAPAM; hPCIF1; C20orf67; PPP1R121; PCIF1


a 400-999-6126 x cn.market@abclonal.com.cn y www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from HeLa cells, using PCIF1 Rabbit pAb (A18347) at 1:2000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20s.