

Bcl-2(Isoform Beta) Rabbit pAb

Catalog No.: A18415 **1 Publications**

Basic Information

Observed MW

22kDa

Calculated MW

26kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human, Rat

Background

This gene encodes a member of the B cell lymphoma 2 protein family. Members of this family regulate cell death in multiple cell types and can have either proapoptotic or antiapoptotic activities. The protein encoded by this gene inhibits mitochondrial-mediated apoptosis. This protein is an integral outer mitochondrial membrane protein that functions as part of signaling pathway that controls mitochondrial permeability in response to apoptotic stimuli. This protein may also play a role in neuron cell survival and autophagy. Abnormal expression and chromosomal translocations of this gene are associated with cancer progression in numerous tissues. Alternate splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

12043

Swiss Prot

P10417

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 150-199 of mouse Bcl2(Isoform Beta) (NP_803129.2).

Synonyms

Bcl-2; C430015F12Rik; D630044D05Rik; D830018M01Rik; Bcl2(Isoform Beta)

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from Rat liver, using Bcl2(Isoform Beta) Rabbit pAb (A18415) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.