

# Pan Cadherin Rabbit pAb

Catalog No.: A18682 **1 Publications**

## Basic Information

### Observed MW

135kDa

### Calculated MW

### Category

Primary antibody

### Applications

ELISA, WB, IP

### Cross-Reactivity

Human, Mouse, Rat

## Background

Cadherin is one of a class of integral-membrane glycoproteins that are involved in cell to cell attachment for preserving the integrity of all solid tissues. Cadherins have three major regions: the  $\text{Ca}^{2+}$ -dependent extracellular region that mediates adhesion (cadherin to cadherin) for cell to cell binding; the transmembrane region; and the cytoplasmic region that extends into the cell and interacts with catenins, which in turn are linked to the actin of the cytoskeleton. Cadherins are differentially expressed during development and in adult organs. Since many cell types express multiple cadherin subclasses simultaneously (the combination differs with cell type), it can be inferred that the adhesion properties of individual cells are thus governed by varying the combinations of cadherins. Altered expression of cadherins are involved in invasion and metastasis of tumour cells. The classical cadherins (e.g. E-, N-, and P-cadherins) are the most common family members. E-cadherin (also known as uvomorulin) is concentrated in the belt desmosome in epithelial cells; N-cadherin is found in nerve, muscle, and lens cells and helps maintain the integrity of neuronal aggregates; P-cadherin is expressed in placental and epidermal cells.

## Recommended Dilutions

**WB** 1:500 - 1:2000**IP** 0.5 $\mu\text{g}$ -4 $\mu\text{g}$  antibody for  
200 $\mu\text{g}$ -400 $\mu\text{g}$  extracts of  
whole cells

## Immunogen Information

### Gene ID

999/1000/1001/1002

### Swiss Prot

P12830/P19022/P22223/P55283

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 800-882 of human pan-cadherin (NP\_004351.1).

### Synonyms

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at  $-20^{\circ}\text{C}$ . Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Immunoprecipitation analysis of 900 µg extracts of PC-3 cells using 3 µg Pan Cadherin antibody (A18682). Western blot was performed from the immunoprecipitate using Pan Cadherin antibody (A18682) at a dilution of 1:3000.



Western blot analysis of various lysates using Pan Cadherin Rabbit pAb (A18682) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.