

Pseudouridine / 5-ribosyluracil Rabbit pAb

Catalog No.: A18872

Basic Information

Observed MW

Refer to figures

Calculated MW**Category**

Primary antibody

Applications

DB, ELISA

Cross-Reactivity

Species independent

Background

Pseudouridine is the most abundant post-transcriptional RNA modification, which presents mostly in non-coding RNAs such as tRNA, rRNA, snRNA and snoRNA. Pseudouridine impacts various aspects of RNA biology, conferring distinct structural and functional properties to the RNA molecules that it decorates. It is also reported that pseudouridine has been found in mRNA in both yeast and human by sequencing analysis. Replacing uridine to pseudouridine enhances structural stability of RNA, and is expected to affect rRNA processing, translation and pre-mRNA splicing. Also, aberrant pseudouridylation contributes to a variety of human diseases, including cancer and genetic disorders. Dysregulation of the pseudouridine epitranscriptome can arise from mutations and abnormal expression of pseudouridylation machinery, impacting protein translation and other cellular processes.

Recommended Dilutions

DB 1:3000 - 1:10000**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID**Swiss Prot****Immunogen**

Chemical compounds corresponding to Pseudouridine / 5-ribosyluracil.

Synonyms

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Dot-blot analysis of all sorts of chemical compounds using Pseudouridine / 5-ribosyluracil antibody (A18872) at 1:5000/1:10000 dilution