[KO Validated] CD44 Rabbit mAb



www.abclonal.com

Catalog No.: A19020 KO Validated Recombinant

19 Publications

Basic Information

Observed MW

80-95 kDa

Calculated MW

82kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,FC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC52411

Background

The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis.

Recommended Dilutions

WB 1:10000 - 1:40000

1:1000 - 1:5000 **IHC-P**

1:200-1:2000 IF/ICC

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 960 P16070

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; CDW44; CSPG8; H-CAM; HCELL; ECM-III; HUTCH-1; HUTCH-I; ECMR-III; Hermes-1; CD44

Contact

| a | 400-999-6126 |
|---------------------|---------------------------|
| \bowtie | cn.market@abclonal.com.cn |
| $\overline{\alpha}$ | www.abclonal.com.cn |


Product Information

Source Isotype **Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using [KO Validated] CD44 Rabbit mAb (A19020) at 1:20000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.




Western blot analysis of lysates from wild type (WT) and CD44 knockout (KO) HeLa cells using [KO Validated] CD44 Rabbit mAb at 1:24000 dilution incubated overnight at 4° C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) ($\overline{A}S014$) at 1:10000 dilution. Lysates/proteins: 25 μg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.




Immunohistochemistry analysis of paraffinembedded Human liver cancer using [KO Validated] CD44 Rabbit mAb (A19020) at dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.





Immunohistochemistry analysis of paraffinembedded Human skin using [KO Validated] CD44 Rabbit mAb (A19020) at dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of A-431 cells using [KO Validated] CD44 Rabbit mAb (A19020, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with $\alpha\text{-Tubulin}$ Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.







Flow cytometry: 1X10^6 Human PBMC were surface-stained with Rabbit IgG isotype control (AC042,2 µg/mL,Ieft) or [KO Validated] CD44 Rabbit mAb (A19020,2.5 µg/mL,right), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Cells in the lymphocyte gate were used for analysis.