

[KD Validated] CDX2 Rabbit mAb

Catalog No.: A19030 **Recombinant** **2 Publications**

Basic Information

Observed MW

38kDa

Calculated MW

34kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0450

Background

This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth and differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis.

Recommended Dilutions

WB	1:1000 - 1:2000
IHC-P	1:500 - 1:2000
IF/ICC	1:100 - 1:1000
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Immunogen Information

Gene ID

1045

Swiss Prot

Q99626

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human CDX2 (Q99626).

Synonyms

CDX3; CDX-3; CDX2/AS; X2

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Immunoprecipitation analysis of 600 μ g extracts of 293T cells using 3 μ g [KD Validated] CDX2 Rabbit mAb (A19030). Western blot was performed from the immunoprecipitate using CDX2 antibody (A19030) at a dilution of 1:500.



Western blot analysis of lysates from wild type (WT) and CDX2 knockdown (KD) 293T cells, using [KD Validated] CDX2 Rabbit mAb (A19030) at 1:500 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of various lysates using [KD Validated] CDX2 Rabbit mAb (A19030) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Exposure time: 3min.


Validation Data



Confocal imaging of Human colon using [KD Validated] CDX2 Rabbit mAb (A19030,dilution 1:100)(Red). DAPI was used for nuclear staining (blue). Objective: 40x.



Immunohistochemistry analysis of paraffin-embedded Human appendix using [KD Validated] CDX2 Rabbit mAb (A19030) at dilution of 1:800 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using [KD Validated] CDX2 Rabbit mAb (A19030) at dilution of 1:800 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.