

# NeuN Rabbit mAb

**Catalog No.: A19086**   **Recombinant**   **18 Publications**

## Basic Information

**Observed MW**

46-55kDa

**Calculated MW**

34kDa

**Category**

Primary antibody

**Applications**

WB,IF-F,IF-P,IHC-P,mIHC,ELISA

**Cross-Reactivity**

Human, Mouse, Rat

**CloneNo number**

ARC0202

## Background

This gene encodes a member of the RNA-binding FOX protein family which is involved in the regulation of alternative splicing of pre-mRNA. The protein has an N-terminal proline-rich region, an RNA recognition motif (RRM) domain, and a C-terminal alanine-rich region. This gene produces the neuronal nuclei (NeuN) antigen that has been widely used as a marker for post-mitotic neurons. This gene has its highest expression in the central nervous system and plays a prominent role in neural tissue development and regulation of adult brain function. Mutations in this gene have been associated with numerous neurological disorders. Alternative splicing of this gene results in multiple transcript variants encoding distinct isoforms.

## Recommended Dilutions

**WB**      1:5000 - 1:20000

**IF-F**      1:200 - 1:800

**IF-P**      1:200 - 1:800

**IHC-P**      1:1000-1:4000

**mIHC**      1:1000-1:4000

**ELISA**      Recommended starting concentration is 1 µg/mL.  
 Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID**

146713

**Swiss Prot**

A6NFn3

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

FOX3; NEUN; FOX-3; HRNBP3; NeuN

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**

Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



## Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)



## Validation Data



Confocal imaging of paraffin-embedded Human brain tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.


Confocal imaging of paraffin-embedded Mouse brain tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Confocal imaging of paraffin-embedded Rat brain tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



The multiplex IHC analysis on paraffin-embedded Mouse brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : NeuN Rabbit mAb (A19086, 1:2000) with TSA-TYR-520 (Green), GFAP Rabbit mAb (A19058, 1:500) with TSA-TYR-570 (Red), and TMEM119 Rabbit mAb (A27143, 1:600) with TSA-TYR-690 (Yellow). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M


The multiplex IHC analysis on paraffin-embedded Rat brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : NeuN Rabbit mAb (A19086, 1:2000) with TSA-TYR-520 (Green), GFAP Rabbit mAb (A19058, 1:500) with TSA-TYR-570 (Red), and TMEM119 Rabbit mAb (A27143, 1:600) with TSA-TYR-690 (Yellow). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M



Confocal imaging of frozen sections Mouse brain(Frozen section) tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.


## Validation Data

citrate buffer at pH 6.0. The analysis was completed using a 20x objective lens.




Immunohistochemistry analysis of paraffin-embedded Human brain tissue using NeuN Rabbit mAb (A19086) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

citrate buffer at pH 6.0. The analysis was completed using a 20x objective lens.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using NeuN Rabbit mAb (A19086) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using NeuN Rabbit mAb (A19086) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.